



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number: **0 477 961 A2**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **91116485.3**

(22) Date of filing: **26.09.91**

(51) Int. Cl.<sup>5</sup>: **C12N 15/53, C12N 15/62,  
C12N 9/02, C12N 15/81,  
C12N 1/19, C12P 7/22**

(30) Priority: **26.09.90 JP 258262/90**  
**24.05.91 JP 120123/91**

(43) Date of publication of application:  
**01.04.92 Bulletin 92/14**

(84) Designated Contracting States:  
**CH DE GB LI**

(71) Applicant: **SUMITOMO CHEMICAL COMPANY,  
LIMITED**  
**5-33, Kitahama 4-chome Chuo-ku**  
**Osaka(JP)**

(72) Inventor: **Akiyoshi, Megumi**  
**4-1-19-205, Nakanokita**  
**Itami-shi Hyogo 664(JP)**  
Inventor: **Yabusaki, Yoshiyasu**  
**6-2-1, Minatojimanakamachi Chuo-ku**  
**Kobe-shi Hyogo 650(JP)**  
Inventor: **Sakaki, Toshiyuki**  
**1-30-68, Higashitoyonakacho**  
**Toyonaka-shi Osaka 560(JP)**  
Inventor: **Ohkawa, Hideo**  
**14-3-403, Asahigaokacho**  
**Ashiya-shi Hyogo 659(JP)**

(74) Representative: **Vossius & Partner**  
**Siebertstrasse 4 P.O. Box 86 07 67**  
**W-8000 München 86(DE)**

(54) **Mitochondrial P450.**

(57) The expression of mitochondrial P450 in yeast is disclosed. The mitochondrial P450 includes a chimeric P450 in which a signal sequence present at the N-terminus of a mammalian mitochondrial P450 has been substituted by a signal of a yeast mitochondrial protein and a chimeric P450 in which the latter signal has been further substituted by a targeting signal sequence to microsome.

An expression plasmid for producing the enzyme in a large scale, a recombinant yeast strain carrying the expression plasmid, a process for producing the enzyme by the use of the recombinant yeast, and a process for producing 5 $\beta$ -cholestane-3 $\alpha$ , 7 $\alpha$ , 12 $\alpha$ , 27-tetrol, 25-hydroxyvitamin D<sub>3</sub> and 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> are also disclosed.

EP 0 477 961 A2

The present invention relates to the expression of mitochondrial P450 in yeast. The mitochondrial P450 includes a chimeric P450 in which a signal sequence present at the N-terminus of a mammalian mitochondrial P450 has been substituted by a signal of a yeast mitochondrial protein and a chimeric P450 in which the latter signal has been further substituted by a targeting signal sequence to microsome. More specifically, the present invention relates to genes which encode a rat liver mitochondria P450<sub>c25</sub> precursor, a chimeric P450 in which a matured P450<sub>c25</sub> is connected with 29 amino acid residues which is believed to be the mitochondrial targeting signal of the yeast cytochrome c oxidase subunit IV (COXIV), and a chimeric P450 in which a signal sequence comprising 15 amino acid residues at the N-terminus of bovine adrenal P450<sub>17α</sub> which is believed to be the microsomal targeting signal is followed by a matured P450<sub>c25</sub>, respectively.

The present invention further relates to an expression plasmid for producing the enzyme in a large scale, a recombinant yeast strain carrying the expression plasmid, a process for producing the enzyme by the use of the recombinant yeast, and a process for producing 5β-cholestane-3α,7α,12α,27-tetrol (referred to hereinafter as TeHC), 25-hydroxyvitamin D<sub>3</sub> and 1α,25-dihydroxyvitamin D<sub>3</sub>.

The term chimeric P450 used in this specification means hereinafter P450 comprising a foreign N-terminal targeting signal sequence and a matured mitochondrial P450 at C-terminus.

P450 is a hemoprotein existing widely in biological fields from microorganisms to mammals and catalyzes as a terminal enzyme of electron transport chains monooxygenation toward a variety of lipophilic compounds as substrates.

P450 as the terminal enzyme in electron transport chains has a variety of molecular forms, which exhibit different substrate specificities and thus can catalyze the hydroxylation of a wide variety of lipophilic compounds. P450-dependent electron transport chains in mammals are classified into two groups; microsomal and mitochondrial types.

In microsomes, NADPH-cytochrome P450 reductase (reductase) containing flavin adenine dinucleotide and flavin mononucleotide as cofactors in the molecule transfers electrons from NADPH to P450. In mitochondria, NADPH-ferredoxin reductase containing flavin adenine dinucleotide as a cofactor in the molecule and ferredoxin containing non-heme iron as a cofactor in the molecule transfer electrons from NADPH to P450.

The present inventors have already succeeded in industrially useful hydroxylations by producing several microsomal P450s and reductases in yeast and using their recombinant yeast strains.

That is, yeast strains producing the enzymes have been obtained by isolating genes for rat liver P450<sub>c</sub>, bovine adrenal P450<sub>17α</sub> and P450<sub>c21</sub>, respectively, making expression plasmids containing these genes, respectively, and transforming yeast with the expression plasmids (Japanese Patent Kokai (Laid-Open) Nos. 56072/1986, 47380/1989 and 31680/1990).

These yeast strains exhibited monooxygenase activities depending on P450 molecular forms produced therein, respectively.

The present inventors have also succeeded in the expression of the enzyme having P450 reducing ability within yeast by isolating rat liver reductase gene or yeast reductase gene (Japanese Patent Kokai (Laid-Open) Nos. 19085/1987 and 51525/1990).

Moreover, the present inventors have succeeded in creating a yeast strain which produces both P450 and reductase (Japanese Patent Kokai (Laid-Open) No. 104582/1987) and novel monooxygenases having the functions of both enzymes (Japanese Patent Kokai (Laid-Open) Nos. 44888/1988 and 23870/1990, and Japanese patent Application No. 71250/1989).

By these techniques, the present inventors have successfully produced acetaminophene or steroid hormone intermediates useful as medicines with these P450-producing yeast strains.

On the other hand, the mitochondrial P450 participates in many biological reactions which synthesize physiologically important compounds such as active vitamin D<sub>3</sub> and steroid hormones, and thus the creation of a yeast strain producing the mitochondrial P450 molecular form has a high industrial applicability.

However, there has not hitherto been reported the expression of an active mitochondrial P450 in yeast.

Furthermore, two enzymes, ferredoxin and ferredoxin reductase, are required in addition of P450 for the expression of mitochondrial P450-dependent monooxygenase activity.

Thus, the present inventors have tried to create a bioreactor for the expression of the mitochondrial P450 in yeast and of the mitochondrial P450-dependent monooxygenase activity.

Rat liver P450<sub>c25</sub> which is a mitochondrial P450 is synthesized as a precursor comprising 533 amino acids in cytoplasm and then transported to mitochondria. During these processes, 32 amino acids at the amino terminus acting as a mitochondrial targeting signal are removed to yield a matured P450<sub>c25</sub> comprising 501 amino acids (molecular weight: 57 KD) on the mitochondrial membrane.

The present inventors have already found that when a bovine adrenal mitochondrial protein is

expressed in yeast, the substitution of its signal peptide part by that of yeast cytochrome c oxidase subunit IV (referred to hereinafter as COXIV) increases the expression amount remarkably (Japanese Patent Application No. 136496/1990).

Rat liver P450<sub>c25</sub> gene or a gene encoding the chimeric P450 (COX-C25) in which its own signal sequence had been substituted by the COXIV signal sequence was linked to the downstream of the promoter of a yeast alcohol dehydrogenase (ADH) gene for construction of expression plasmids. As a result, yeast strains transformed with these expression plasmids produced P450s containing heme.

It was also found that the P450<sub>c25</sub> and the chimeric P450 (COX-C25) thus produced were transported to the mitochondria in yeast and processed into the matured forms. Furthermore, the P450<sub>c25</sub> and the chimeric P450 (COX-C25) partially purified from these transformed yeast strains successfully produced TeHC or 1 $\alpha$ , 25-dihydroxyvitamin D<sub>3</sub> from THC or 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> by coupling with bovine adrenal adrenodoxin (referred to hereinafter as ADX) and bovine adrenal NADPH-adrenodoxin reductase (referred to hereinafter as ADR).

Furthermore, the present inventors have constructed a plasmid for expressing the chimeric P450 (17 $\alpha$ -c25) in which the mitochondrial targeting signal substituted by the N-terminal 15 amino acids derived from bovine adrenal P450<sub>17 $\alpha$</sub> , which is believed to be a microsomal targeting signal of P450<sub>17 $\alpha$</sub>  in order to enhance the production of the enzyme inherently existing in mitochondria by exchanging its localization into microsomes.

The present inventors have also constructed a plasmid which express simultaneously the chimeric P450 (17 $\alpha$ -c25), the matured ADX and the matured ADR for the purpose of constructing a novel electron transport chain with the chimeric P450 (17 $\alpha$ -c25) on microsomal membrane and ADX and ADR in cytoplasm and exhibiting the monooxygenase activity. In this connection, these three enzymes were expressed with different promoters and terminators, because the use of the same promoter and the same terminator would cause recombination in the plasmid during the cultivation of the recombinant yeast.

It was found that all of the P450<sub>c25</sub>, the chimeric P450 (COX-c25) and the chimeric P450 (17 $\alpha$ -c25) expressed in yeast in the present invention exhibited hydroxylation activities to THC, vitamin D<sub>3</sub> and 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> and thus were active forms. Thus, the yeast expression system is useful for expressing the mitochondrial P450.

Particularly, the chimeric P450 (17 $\alpha$ -c25) produces a large amount of the enzyme and has a high activity.

While P450<sub>c25</sub> and the chimeric P450 (COX-c25) were expressed in mitochondria in yeast, any chimeric P450 (17 $\alpha$ -c25) expressed existed in microsomal fraction. Thus, the N-terminal 15 amino acid residues derived from P450<sub>17 $\alpha$</sub>  played a role of a targeting signal of the chimeric P450 (17 $\alpha$ -c25) to microsomal membrane in yeast, and the P450<sub>c25</sub> which is inherently mitochondrial membrane protein was successfully expressed in yeast microsome.

The production amount of the chimeric P450 (17 $\alpha$ -c25) was about 2 x 10<sup>5</sup> molecules/cell, and thus the substitution of the signal successfully increased the production amount to about 4 times. Moreover, the microsomal fraction exhibited high hydroxylation activities to THC, vitamin D<sub>3</sub> and 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> by adding ADX and ADR to the fraction.

Particularly, the hydroxylation activity to 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> of the chimeric P450(17 $\alpha$ -c25) in the microsomal fraction was by far higher than that of the original matured P450<sub>c25</sub> comprising 501 amino acids in the mitochondrial fraction.

The present inventors has also succeeded in construction of a novel electron transport chain from NADPH to the chimeric P450 (17 $\alpha$ -c25) through ADX and ADR on yeast microsomal membrane, as the simultaneous expression strain of chimeric P450 (17 $\alpha$ -c25), ADX and ADR exhibits monooxygenase activity.

Therefore, the strain can be used as a simple bioreactor in which the hydroxylation of steroid compounds or an active vitamin D<sub>3</sub> is performed in one step.

Particularly, the strain can easily recover reaction products by adding a substrate to the culture of the microorganism and extracting with an organic solvent or the like after a certain period of time, so that it is useful as a bioreactor.

The figures show:

Fig. 1 shows the constructing process of the expression plasmid pAC25 of the present invention.

Fig. 2 shows a map of pAC25.

Fig. 3 shows the constructing process of the expression plasmid pAC253 of the present invention, in which the restriction enzyme sites represent the followings: Ec: EcoRI, Nc: NcoI, Sc: SacI, Hd: HindIII, Nt: NotI, Ps: PstI, and in the drawing,  $\square$  represents rat liver P450<sub>c25</sub> coding region,



5 represents yeast cytochrome c oxidase subunit IV coding region, and P and T represent ADH promoter and terminator, respectively.

Fig. 4 is a map of PAC253.

Fig. 5 is the explanatory drawing of the constructing process of the expression plasmid of the present invention pAMS25, in which the restriction enzyme cutting sites are represented as follows: E: EcoRI, Nc:  
10 NcoI, S: SacI, H: HindIII, P: PstI, N: NotI, and

Fig. 6 is the explanatory drawing of the constructing process of the expression plasmid of the present invention pRXMS25, in which the restriction enzyme cutting sites are represented as follows: H: HindIII and N: NotI.

The present invention will be explained in detail below.

15 cDNA encoding the rat liver P450<sub>c25</sub> used in the present invention is well known and can be isolated by the conventional procedures. The signal sequence portion to mitochondria can be substituted by a signal sequence of the other mitochondrial protein such as COXIV or the like. In this case, P450 is transported into mitochondria. On the other hand, if this signal sequence is substituted by a signal sequence to microsome, P450 is transported into microsome. As the transfer signal to microsome, not only bovine adrenal P450<sub>17α</sub>  
20 but also a transfer signal of the other microsomal P450 can be used.

Furthermore, as the matured mitochondrial P450 existing at the C-terminus of the chimeric P450 of the present invention, not only the rat liver P450<sub>c25</sub> but also the other mitochondrial P450 can be used.

cDNA encoding COXIV and bovine adrenal P450<sub>17α</sub> used in the present invention is well known and can be isolated by the conventional procedures.

25 For example, the expression plasmid for expressing the chimeric P450 (17α-c25) can be constructed by synthesizing cDNA corresponding the N-terminal amino acid sequence of bovine adrenal P450<sub>17α</sub>, linking to the cDNA region encoding the matured rat liver P450<sub>c25</sub>, and inserting the gene into the yeast expression vector pAAH5 which retains the promoter and the terminator of the yeast alcohol dehydrogenase I (referred to hereinafter as ADH) gene (Methods in Enzymology, 101, part C, 192-201).

30 The plasmid for simultaneously expressing the three enzymes can be constructed by inserting the expression units of the matured ADR and the matured ADX into the expression plasmid for the chimeric P450 (17α-c25). In this connection, the promoter and the terminator are not limited to ADH and may be the promoters and the terminators which function efficiently in yeast. The promoter and the terminator for expressing simultaneously the three enzymes are not the same and preferably different from each other,  
35 because recombination on plasmid will occur only with a low probability upon cultivation. The expression amount of the enzyme is scarcely affected by the position or the direction of the three enzymes to be inserted into the expression unit.

Yeast is used as a host, and the yeast *Saccharomyces cerevisiae* strain AH22 is preferred.

Transformation of yeast as a host by the expression plasmid can be conducted by the well-known  
40 methods such as that with an alkali metal (LiCl) or protoplast. Transformed yeast obtained by the present invention can be cultivated in an ordinary culture medium containing glucose, nitrogen sources or the like by the conventional methods. P450<sub>c25</sub> and the chimeric P450 can be produced by cultivating the transformed yeast strain thus obtained.

The present invention is described in detail with reference to Examples. However, it goes without saying  
45 that the present invention is not limited to Examples.

#### Example 1 Construction of expression plasmid pAC25

In the following Examples, the reactions such as the cutting of DNA with restriction enzymes, the  
50 dephosphorylation of DNA with alkaline phosphatase, the ligation of DNAs with DNA ligase were conducted generally with a reaction volume of 20 - 200 μl under the reaction conditions described in the instructions appended in the products of the manufacturers (e.g. Takara Shuzo Co., Ltd.) of these enzymes, unless otherwise specified. Rat liver P450<sub>c25</sub> expression plasmid was constructed in such a procedure as shown in Fig. 1.

55 Plasmid pLMT25 containing rat liver P450<sub>c25</sub> gene (Usui et al., (1990) FEBS Lett., 262, 135-138) was doubly digested with restriction enzymes EcoRI and SacI, and the ca. 320 bp EcoRI-SacI fragment encoding the region of the amino terminal side of P450<sub>c25</sub> and the ca. 1580 bp SacI-EcoRI fragment encoding the region of the carboxy terminal side were recovered by the low melting agarose gel

electrophoresis. These fragments were subcloned at the EcoRI and SacI sites of a commercially available vector plasmid pUC19 to give pUC25N and pUC25C, respectively. A fragment of about 2880 bp obtained by the doubly digestion of pUC25N with EcoRI and NcoI, and a synthetic linker LC252:

5' AATTCAAGCTTAAAAAATGGCTCTGTTGAGCCGCATGAGA  
3' GTTCGAATTTTTTTACCGACACAACCTCGGCGTACTCT  
CTGAGATGGGCGCTTCTGGACACTCGTGTGATGGGC 3'  
GACTCTACCCGCGAAGACCTGTGAGCACACTACCCGGTAC 5'

which has EcoRI and NcoI recognition sites at right and left ends, respectively, and HindIII recognition site between the both ends, were ligated for transform *E. coli* strain HB101.

Plasmid DNA was prepared from the transformant thus obtained according to the Birnboim-Doly's method and analyzed by the digestion with restriction enzymes to obtain the aimed plasmid in which the synthetic linker was inserted. Further, the sequence of the synthetic linker part was confirmed by the determination of the base sequence. Thus, the plasmid obtained was named pUC25NH.

The plasmid pUC25C encoding the region of the carboxy terminal side of P450<sub>c25</sub> was digested with NcoI and fill-in ligated with a commercially available HindIII linker to transform *E. coli* strain HB101. Plasmid DNA was prepared from the transformant, and the plasmid in which a HindIII site was created in the 3'-non-coding region of P450<sub>c25</sub> was named pUC25CH.

The fragments of about 270 bp and 1370 bp obtained by the doubly digestion of pUC25NH and pUC25CH with HindIII and SacI, respectively, and a commercially available vector plasmid pUC18 which had been digested with HindIII and treated with alkaline phosphatase were ligated to transform *E. coli* strain HB101. A plasmid DNA was prepared from the transformant, and the plasmid containing both DNA fragments derived from pUC25NH and pUC25CH, respectively, was named pUC25H. pUC25H was digested with Hind III, and a cDNA fragment of ca. 1640 bp encoding the rat liver P450<sub>c25</sub> precursor was prepared. After the fragment and the yeast expression vector pAAH5N which had been digested with HindIII and then treated with alkaline phosphatase (Japanese Patent Application No. 202785/1988) were ligated, *E. coli* strain HB101 was transformed. A plasmid DNA was prepared from the transformant, the DNA structure was confirmed by restriction enzyme digestion and the plasmid in which the rat liver P450<sub>c25</sub> gene had been inserted in the right orientation between the ADH promoter and the terminator was named pAC25.

#### Example 2 Construction of expression plasmid pACC253

Reference is made to Fig. 3.

Plasmid pUC25N containing the gene encoding the region of the amino terminal side of the rat liver P450<sub>c25</sub> was doubly digested with PstI and SacI to recover a fragment of about 150 bp. The fragment, the synthetic linker LC253:

5' AGCTTAAAAAATGCTTTTCACTACGTCAATCGATAAGATTTTTCAAGCCAGCCACAAGAAC  
3' ATTTTTTTACGAAAGTGATGCAGTTAGCTATTCTAAAAAGTTCGGTCGGTGTCTTG  
TTTGTGTAGCTCTAGATATCTGCTTCAGCAAAAACCCGCGATCCCTGCA 3'  
AAACACATCGAGATCTATAGACGAAGTCGTTTTTGGGCGCTAGGG 5'

which has HindIII and PstI recognition sites at right and left ends, respectively, and the HindIII-SacI fragment of a commercially available vector pUC19 were ligated, and *E. coli* strain HB101 was transformed. Plasmid

DNA was prepared from the transformant thus obtained and analyzed by the digestion with restriction enzymes to obtain the aimed plasmid in which the fragment and the synthetic linker were inserted. Further, the sequence of the synthetic linker part was confirmed by the determination of the base sequence. Thus, the plasmid obtained was named pBSCC253. The plasmid and the aforementioned plasmid pUC25CH were  
 5 doubly digested with HindIII and SacI, and a DNA fragment (ca. 260 bp) coding for the signal peptide of COXIV and the amino terminal part of P450<sub>c25</sub> and a DNA fragment (ca. 1370 bp) coding for the carboxy terminal part of P450<sub>c25</sub> were recovered separately. The triple ligation of these fragments and a commercially available vector plasmid pUC18-HindIII fragment was conducted to transform *E. coli* strain HB101. A plasmid DNA was prepared from the transformant and analyzed by the digestion with restriction enzymes to  
 10 obtain a plasmid pUCC253H in which the fragments had been inserted. pUCC253H was digested with HindIII to prepare a modified cDNA fragment (ca.1630 bp) for the rat liver P450<sub>c25</sub>, which was ligated with the yeast expression vector pAAH5N which had been digested with HindIII and then treated with alkaline phosphatase (Japanese Patent Application No. 202785/1988), and then *E. coli* strain HB 101 was transformed. The structure of the plasmid DNA prepared from the transformant was confirmed, and the plasmid  
 15 in which the rat liver P450<sub>c25</sub> modified gene had been inserted in the right orientation between the ADH promoter and the terminator was named pACC253.

#### Example 3 Construction of expression plasmid pAMS25 and simultaneous expression plasmid pRXMS25

20 The chimeric P450 (17 $\alpha$ -c25) expression plasmid pAMS25 was constructed according to Fig. 5.  
 The plasmid pUC25N used in the rat liver P450<sub>c25</sub> expression plasmid pAC25 illustrated in Example 1 was partially digested with PstI and then with EcoRI. Into the DNA fragment obtained was inserted a synthetic linker: sequence

25 **AATTCAAGCT TAAAAAATG TGGCTGCTCC TGGCTGTCTT TCTGCTCACC CTCGCTATT 60**  
**GTTCGA ATTTTTTTTAC ACCGACGAGG ACCGACAGAA AGACGAGTGG GAGCGGATAA**  
 30 **TAGCGATCCC TGCA** 74  
**ATCGCTAGGG**

35 and a HindIII-SacI fragment was prepared from the plasmid thus obtained. Also, a SacI-HindIII fragment was prepared from pUC25CH and doubly inserted into the HindIII site of pUC19 for obtaining a plasmid pUMS25.

Next, an expression plasmid pAMS25 was constructed by inserting the HindIII fragment obtained from  
 40 pUMS25 into the Hind III site of a vector pAAH5N.

An expression plasmid pGX was obtained by inserting a Hind III fragment coding for the matured ADX (described in Japanese Patent Application No. 136496/1990) between the PGK promoter and the terminator of a vector pGAHN (containing the PGK promoter and the terminator in place of the ADH promoter and the terminator of a vector pAAH5N) (described in Japanese Patent Application No. 136496/1990). pPR was also  
 45 prepared by inserting a Hind III fragment coding for the matured ADR (described in Japanese Patent Kokai No. 112986/1988) between the GAP promoter and the terminator of a vector pPAHN, which contains the GAP promoter and the terminator (Japanese Patent Kokai (Laid-Open) No. 112986/1988) in place of the ADH promoter and the terminator of the vector pAAH5N). Simultaneous expression plasmid pRX5 for ADR and ADX was obtained by ligating the DNA fragment obtained by partially digesting pPR with NotI and the  
 50 NotI fragment obtained from pGX. Then, pRX5 was partially digested with NotI, ligated with a NotI fragment (3.7 kb) obtained from pAMS25 to obtain the simultaneous expression plasmid pRXMS25 for the chimeric P450<sub>c25</sub>, ADX and ADR (See Fig. 6).

#### Example 4 Transformation of yeast with expression plasmid

55 *Saccharomyces cerevisiae* AH22 (ATCC 38626) was cultivated in 5 ml of a YPD culture medium (1 % yeast extract, 2 % polypeptone, 2 % glucose) at 30 °C for 18 hours, and the yeast cells were collected by centrifuging 1 ml of the culture. The yeast cells were washed with 1 ml of a 0.2 M LiCl solution and then

suspended into 20  $\mu$ l of a 1 M LiCl solution.

To this suspension were added 30  $\mu$ l of a 70 % polyethylene glycol 4000 solution and 10  $\mu$ l of respective expression plasmid solutions (about 1  $\mu$ g DNA), and the mixture was stirred well and incubated at 30°C for 1 hour. Then, 140  $\mu$ l of water was added to the mixture, and the resulting mixture was inoculated on an SD synthetic culture medium plate (2 % glucose, 0.67 % yeast nitrogen base without amino acids, 20  $\mu$ g/ml histidine, 2 % agar) and incubated at 30°C for 3 days to give a transformant in which the plasmid was maintained.

#### Example 5 Production of P450<sub>c25</sub> and chimera P450<sub>c25</sub>

The strains AH22/pAC25, AH22/pACC253, AH22/pAMS25 and AH22/pRXMS25 obtained in Example 4 were cultivated in 300 ml of a synthetic culture medium (8 % glucose, 5.4 % yeast nitrogen base without amino acids, 160  $\mu$ g/ml histidine) to a density about  $2 \times 10^7$  cells/ml, and the recombinant yeast cells were collected, washed with 100 mM potassium phosphate (pH 7.0) and suspended into 2 ml of the same buffer solution. A 1 ml portion of the suspension was dispensed into each of the two cuvettes, carbon monoxide was blown into the sample cuvette, and then 5 - 10 mg of dithionite was added to both of the cuvettes. After sufficiently stirring the mixtures, the differential spectra were measured in the wave length of 400 - 500 nm to calculate the amount of the heme-containing P450 based on  $\Delta\epsilon$  (450 nm - 490 nm) = 91 mM<sup>-1</sup>cm<sup>-1</sup>.

As the result, it was found that the strains AH22/pAC25, AH22/pACC253, AH22/pAMS25 and AH22/pRXMS25 produced a heme-containing P450 in an amount of  $5 \times 10^4$  molecules,  $3 \times 10^4$  molecules,  $2 \times 10^5$  molecules and  $1 \times 10^5$  molecules per cell, respectively. On the other hand, no production of the heme-containing P450 was observed in the control strain AH22/pAAH5.

#### Example 6 Measurement of the hydroxylation activity at 25 position of 1 $\alpha$ -hydroxy-vitamin D<sub>3</sub> and the hydroxylation activity at 27 position of THC

A mitochondrial fraction prepared from the strain AH22/pAC25 and a microsomal fraction prepared from the strain AH22/pAMS25 were used for the measurement of hydroxylation activities. Subcellular fractionation was prepared from a transformant yeast strain according to the following method. About  $4 \times 10^{10}$  cells were collected from a culture containing about  $2 \times 10^7$  cells/ml, suspended in Zymolyase solution [10 mM Tris-HCl (pH 7.5), 2.0 M sorbitol, 0.1 mM dithiothreitol, 0.1 mM EDTA, 0.3 mg/ml Zymolyase 100T] and incubated at 30°C for 1 hour to prepare spheroplasts. The spheroplasts were suspended in a sonication buffer (10 mM Tris-HCl (pH 7.5), 0.65 M sorbitol, 0.1 mM dithiothreitol, 0.1 mM EDTA, 1  $\mu$ g/ml leupeptin, 1  $\mu$ g/ml pepstatin) and homogenized with a Teflon homogenizer to disrupt the cells.

Precipitate 1 was removed by centrifugation at 3000 x g for 5 minutes to give supernatant 1'. Precipitate 1 was resuspended in a sonication buffer, homogenized again, centrifuged at 3000 x g for 5 minutes to give precipitate 1 (undisrupted cells and nucleus fractions) and supernatant 1''. Supernatant 1 comprising supernatant 1' and supernatant 1'' was further centrifuged at 10,000 x g for 20 minutes to give precipitate 2 (mitochondrial fraction) and supernatant 2. Supernatant 2 was further centrifuged at 120,000 x g for 70 minutes to give precipitate 3 (microsomal fraction) and supernatant 3 (soluble fraction).

Reaction systems (2 ml) shown in the following but NADPH were mixed and preincubated at 37°C for 5 minutes, and reaction was initiated by the addition of NADPH. After 5 and 10 minutes, 0.5 ml portions of reaction solutions were taken, respectively, and 5 ml of benzene was added to the removed samples. The benzene layer obtained by centrifugation was dried completely and analyzed by HPLC under the condition listed below.

#### [Reaction system]

1. Mitochondrial fraction: strain AH22/pAC25 (containing 0.1 nmol P450<sub>c25</sub>),
2. Microsomal fraction: strain AH22/pAMS25 (containing 0.8 nmol chimeric P450 (17 $\alpha$ -c25),
3. Bovine adrenal adrenodoxin reductase 0.8 nmol,
4. Bovine adrenal adrenodoxin 8.0 nmol,
5. Substrate final concentration 200  $\mu$ M,
6. NADPH final concentration 1.0-2.0 mM,
7. Tris-HCl (pH 7.8) 100.0 mM,
8. EDTA 0.5-1.0 mM.

#### [HPLC analysis condition]

Column:  $\mu$ Bondapak C18 ( $\phi$ 4 x 300 mm),  
 Detection: A265 (1 $\alpha$ -hydroxyvitamin D<sub>3</sub>), Radioactive detector (<sup>3</sup>H-THC),  
 Flow rate: 1.0 ml/min,  
 Temperature: 50 °C,

5

## Elution condition

0 - 5 min: 80 % acetonitrile - 20 % water solution  
 5 - 15 min: linear concentration gradient of 80 % - 100 % acetonitrile,  
 10 15 - 25 min: 100 % acetonitrile

When 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> was used as a substrate, the 25-hydroxylated product were observed in the mitochondrial fraction of the strain AH22/pAC25 and the microsomal fraction of the strain AH22/pAMS25. On the other hand, no 25-hydroxylated products were observed in the control strain AH22/pAAH5. Thus, it was suggested that the hydroxylation activities are dependent on P450<sub>c25</sub> and the chimeric P450 (17 $\alpha$ -c25).

15 In addition, no activity was detected in the absence of either NADPH, ADX or ADR, so that the hydroxylation activities were believed to be exhibited with the construction of the electron transfer chain, NADPH  $\rightarrow$  ADR  $\rightarrow$  P450<sub>c25</sub> or chimeric P450 (17 $\alpha$ -c25), which was formed on the addition of NADPH, ADX and ADR.

In the microsomal fraction of the strain AH22/pAMS25, the calculated turnover number was 7.4 mol/mol P450/min, and thus it was found that the rate was about 50 times higher than the calculated turnover number obtained in the mitochondrial fraction of the strain AH22/pAC25 of 0.14 mol/mol P450/min. When [<sup>3</sup>H]-THC was used as a substrate, the transformation of THC to the 27-hydroxylated product (TeHC) was observed in the mitochondrial fraction of the strain AH22/pAC25 and the microsomal fraction of the strain AH22/pAMS25. No activity was observed in the control strain AH22/pAAH5. Thus, it was suggested that the activity was dependent on P450<sub>c25</sub> and chimeric P450 (17 $\alpha$ -c25). The turnover numbers calculated in the mitochondrial fraction of the strain AH22/pAC25 and the microsomal fraction of the strain AH22/pAMS25 were 20 mol/mol P450/min and 23 mol/mol P450/min, respectively.

#### 30 Example 7 Measurement of the 27-hydroxylation activity of THC in the triple simultaneous expression strain AH22/pRXMS25

To the culture of the strain AH22/pRXMS25 (1.6 x 10<sup>7</sup> cells/ml) was added [<sup>3</sup>H]-THC so that the final concentration is 10  $\mu$ M, and the incubation was continued. A 1 ml portion of the culture after 14 hour cultivation was taken out and extracted with 2 ml of dichloromethane, and the dichloromethane layer was evaporated to dryness, dissolved in 80  $\mu$ l of acetonitrile, of which a 50 $\mu$ l portion was analyzed by HPLC under the same condition as described in Example 6. As a result, it was found that 19 % of the substrate THC added had been transformed into TeHC. Such transformation was not observed in the control strain AH22/pAAH5, and thus it was found that the chimeric P450<sub>c25</sub>, ADX and ADR constructed an electron transport chain and exhibited a monooxygenase activity in the strain AH22/pRXMS25.

40 The transformed yeast provided by the present invention produces P450<sub>c25</sub> containing heme in the molecule and the chimeric P450<sub>c25</sub>.

P450<sub>c25</sub> and the chimeric P450<sub>c25</sub> exhibited 25-hydroxylation activities to vitamin D<sub>3</sub> and 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> and 27-hydroxylation activity to THC by the addition of bovine adrenal ADX and ADR.

45 While P450<sub>c25</sub> is a protein inherently present in mitochondrial inner membrane, conversion from mitochondria to microsome for its localization in yeast cells was succeeded by removing the mitochondrial targeting signal and adding the N-terminal 15 amino acid residues derived from the microsomal P450<sub>17 $\alpha$</sub> , which functions as a targeting signal to microsomal membrane. Such a conversion of the localization of the membrane enzyme has not hitherto been reported in the literatures. While the 15 amino acid residues at the N-terminus of P450<sub>17 $\alpha$</sub>  are highly hydrophobic and is believed to play a role as a targeting signal to microsomal membrane, it has no specificity as a signal and can be replaced with a signal of the other microsomal membrane proteins.

50 The production amount of P450<sub>c25</sub> per yeast cell could be increased to about 4 times and the 25-hydroxylation activity of 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> per P450 molecule could be also increased to about 50 times by changing mitochondria into microsome for the localization of membrane enzyme based on the substitution of the signal. Thus, the ability of the 25-hydroxylation of 1 $\alpha$ -hydroxyvitamin D<sub>3</sub> per cell was assumed to be raised to about 200 times. It is possible to produce 1 $\alpha$ , 25-dihydroxyvitamin D<sub>3</sub> which plays a role for regulating calcium metabolism, stimulating the differentiation of cells and controlling cellular immune systems and is useful for the treatment of osteoporosis, chronic renal insufficiency, vitamin D



resistant rickets or the like by using P450<sub>c25</sub> or the chimeric P450<sub>c25</sub> as a bioreactor.

In addition, the use of the cells of the strain which expresses simultaneously chimeric P450<sub>c25</sub>, ADX and ADR as a bioreactor is a useful means for simplifying the currently used complex method for chemical synthesis of an active vitamin D<sub>3</sub>.

- 5 Further, P450<sub>c25</sub> produced in yeast can be easily purified, and an antibody against P450<sub>c25</sub> can be used as a diagnostic agent of cerebrotendinous xanthomatosis. It has been definitely shown that the patients of cerebrotendinous xanthomatosis are congenitally deficient in the enzyme. The disease is characterized in xanthomatosis or the neuropathy of animus, and xanthomatosis is generated at Achilles tendon, lung and brain of the patient resulting in serious symptoms such as cataract, lowering of intelligence or the like. It is  
10 known that the cholestanol level in the serum of the patients is in a proportion of 10 - 100 to the normal level, and diagnostic methods utilizing this phenomenon or methods of measuring the activity of the enzyme with THC labelled with a radioisotope are currently used. The production of P450<sub>c25</sub> in a large scale has become possible by cultivating the strain of the transformed yeast obtained by the present invention, so that the preparation of an antibody against the isolated and purified enzyme makes it possible to detect  
15 P450<sub>c25</sub> at a high sensitivity and is believed to be effective for the diagnosis of cerebrotendinous xanthomatosis.

20

25

30

35

40

45

50

55

## SEQUENCE LISTING

5

1. SEQ ID NO: 1:

SEQUENCE TYPE: nucleic acid  
 SEQUENCE LENGTH: 110  
 STRANDEDNESS: double  
 TOPOLOGY: straight  
 IMMEDIATE EXPERI-  
 15 MENTAL SOURCE: synthetic linker

5' AGCTTAAAAAATGCTTTCACTACGTCAATCGATAAGATTTT  
 20 3' ATTTTITACGAAAGTCATGCAGTTAGCTATTCTAAAA  
 TCAAGCCAGCCACAAGAACTTTGTGTAGCTCTAGATATCTGC  
 25 AGTTCGGTCGGTCTTCTTCAAACACATCGAGATCTATAGACG  
 TTCAGCAAAAACCCGGCATCCCTGCA 3'  
 30 AAGTCGTTTTTGGGGCTAGGG 5'

35

2. SEQ ID NO: 2:

40 SEQUENCE TYPE: nucleic acid  
 SEQUENCE LENGTH: 74  
 STRANDEDNESS: double  
 TOPOLOGY: straight  
 45 IMMEDIATE EXPERI-  
 MENTAL SOURCE: synthetic linker

50 AATTCAAGCT TAAAAAATG TGGCTGCTCC TGGCTGTCTT TCTGCTCACC CTCGCCTATT 60  
 GTTCGA ATTTTTTAC ACCGACGAGG ACCGACAGAA AGACGAGTGG GAGCGGATAA  
 TAGCGATCCC TGCA 74  
 55 ATCGCTAGGG

## Claims

- 5 1. A plasmid for expressing a mitochondrial P450 or a chimeric mitochondrial P450 in yeast.
2. An expression plasmid according to claim 1, wherein said mitochondrial P450 is a rat liver P450<sub>c25</sub>.
3. An expression plasmid for a mitochondrial P450 or a chimeric mitochondrial P450 in which a targeting  
10 signal at an N-terminus is substituted by a targeting signal of a mitochondrial protein.
4. A plasmid according to claim 2, wherein a targeting signal at an N-terminus is substituted by a targeting signal of a mitochondrial protein.
- 15 5. An expression plasmid for a mitochondrial P450 or a chimeric mitochondrial P450 in which a targeting signal at an N-terminus is substituted by a targeting signal of a microsomal protein.
6. A plasmid according to claim 2, wherein a mitochondrial targeting signal at an N-terminus is substituted by a targeting signal of a microsomal protein.
- 20 7. An expression plasmid according to any one of claims 3 and 4, wherein a mitochondrial targeting signal at an N-terminus is that for yeast mitochondrial COXIV.
8. A plasmid according to any one of claims 5 and 6, wherein a microsomal targeting signal at an N-terminus is the N-terminal 15 amino acids of a bovine adrenal P450<sub>17 $\alpha$</sub> .
- 25 9. A plasmid according to any one of claims 1 to 8 containing a promoter of a yeast alcohol dehydrogenase.
- 30 10. An expression plasmid pAC25 represented by Fig. 2.
11. An expression plasmid pACC253 represented by Fig. 4.
12. An expression plasmid pAMS25 represented by Fig. 5.
- 35 13. A yeast strain carrying an expression plasmid according to any one of claims 1 to 12.
14. A process for producing a mitochondrial P450 or a chimeric mitochondrial P450, in which a yeast strain according to claim 13 is cultivated.
- 40 15. A process for producing 5 $\beta$ -cholestane-3 $\alpha$ , 7 $\alpha$ , 12 $\alpha$ , 27-tetrol, 25-hydroxyvitamin D<sub>3</sub> and 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>, characterized in that a hydroxy group is introduced into the 27-position of 5 $\beta$ -cholestane-3 $\alpha$ ,7 $\alpha$ ,12 $\alpha$ -triol and the 25-position of vitamin D<sub>3</sub> and 1 $\alpha$ -hydroxyvitamin D<sub>3</sub>, respectively.
- 45 16. A plasmid for simultaneously expressing a mitochondrial P450 or a chimeric mitochondrial P450, ferredoxin and ferredoxin reductase.
17. A plasmid according to claim 16, wherein said mitochondrial P450 is a rat liver P450<sub>c25</sub>.
- 50 18. A plasmid according to any one of claims 16 and 17, wherein a targeting signal at an N-terminus of said mitochondrial P450 or said chimeric mitochondrial P450 is substituted by a targeting signal of a mitochondrial protein.
19. A plasmid according to any one of claims 16 to 18, wherein said mitochondrial targeting signal at the  
55 N-terminus is that of a yeast mitochondrial COXIV.
20. A plasmid according to any one of claims 16 and 17, wherein a targeting signal at an N-terminus of said mitochondrial P450 or said chimeric mitochondrial P450 is substituted by a targeting signal of a

microsomal protein.

21. A plasmid according to any one of claims 16, 17 and 20, wherein a microsomal targeting signal at an N-terminus is the N-terminal 15 amino acids of a bovine adrenal P450<sub>17 $\alpha$</sub> .

5

22. A plasmid according to any one of claims 16 to 21, wherein said ferredoxin and said ferredoxin reductase are derived from a bovine adrenal gland.

23. An expression plasmid pRXMS25 represented by Fig. 6.

10

24. A yeast strain carrying a plasmid according to any one of claims 16 to 23.

25. A yeast strain carrying at least one plasmid according to any one of claims 1 to 12 and 16 to 23.

15

20

25

30

35

40

45

50

55

FIG. 1(A)




FIG. 1(B)




FIG. 2




FIG. 4




FIG. 3






FIG. 5





FIG. 6





Europäisches Patentamt  
European Patent Office  
Office européen des brevets



Publication number: **0 477 961 A3**

**EUROPEAN PATENT APPLICATION**

Application number: **91116485.3**

Int. Cl.<sup>5</sup>: **C12N 15/53, C12N 15/62,  
C12N 9/02, C12N 15/81,  
C12N 1/19, C12P 7/22**

Date of filing: **26.09.91**

Priority: **26.09.90 JP 258262/90  
24.05.91 JP 120123/91**

Date of publication of application:  
**01.04.92 Bulletin 92/14**

Designated Contracting States:  
**CH DE GB LI**

Date of deferred publication of the search report:  
**29.04.92 Bulletin 92/18**

Applicant: **SUMITOMO CHEMICAL COMPANY,  
LIMITED  
5-33, Kitahama 4-chome Chuo-ku  
Osaka(JP)**

Inventor: **Akiyoshi, Megumi  
4-1-19-205, Nakanokita  
Itami-shi Hyogo 664(JP)  
Inventor: Yabusaki, Yoshiyasu  
6-2-1, Minatojimanakamachi Chuo-ku  
Kobe-shi Hyogo 650(JP)  
Inventor: Sakaki, Toshiyuki  
1-30-68, Higashitoyonakacho  
Toyonaka-shi Osaka 560(JP)  
Inventor: Ohkawa, Hideo  
14-3-403, Asahigaokacho  
Ashiya-shi Hyogo 659(JP)**

Representative: **Vossius & Partner  
Siebertstrasse 4 P.O. Box 86 07 67  
W-8000 München 86(DE)**

**Mitochondrial P450.**

The expression of mitochondrial P450 in yeast is disclosed. The mitochondrial P450 includes a chimeric P450 in which a signal sequence present at the N-terminus of a mammalian mitochondrial P450 has been substituted by a signal of a yeast mitochondrial protein and a chimeric P450 in which the latter signal has been further substituted by a targeting signal sequence to microsome.

An expression plasmid for producing the enzyme in a large scale, a recombinant yeast strain carrying the expression plasmid, a process for producing the enzyme by the use of the recombinant yeast, and a process for producing 5 $\beta$ -cholestane-3 $\alpha$ , 7 $\alpha$ , 12 $\alpha$ , 27-tetrol, 25-hydroxyvitamin D<sub>3</sub> and 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> are also disclosed.

**EP 0 477 961 A3**

EP 91 11 6485

Page 1

DOCUMENTS CONSIDERED TO BE RELEVANT			Page 1
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	WO-A-8 910 963 (GIST-BROCADES N.V.)	1, 3-5, 7, 13-14, 25	C12N15/53 C12N9/02 C12N15/81 C12N15/62 C12N1/19 C12P33/06
A	* examples 5-6, 8-10, 22-23, 27 and the claims *	16, 22, 24	
	---		
D, Y	FEBS LETTERS. vol. 262, no. 1, March 1990, AMSTERDAM NL pages 135 - 138; EMIKO USUI ET AL: 'Molecular cloning of cDNA for vitamin D3 25-hydroxylase from rat liver mitochondria' * the whole document *	1-2, 8-10, 13-15	
	---		
Y	CHEMICAL ABSTRACTS, vol. 113, no. 1, 2 July 1990, Columbus, Ohio, US; abstract no. 2514W, SHIBATA M. ET AL: 'Genetically engineered P450 monooxygenases: construction of bovine P450c17/yeast reductase fused enzymes' page 258 ; column L ; * abstract * & DNA AND CELL BIOLOGY vol. 9, no. 1, January 1990, NEW YORK; US. pages 27 - 36;	1-2, 8-10, 13-15	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
	---		C12N C12P
Y	US-A-4 766 068 (AGENCY OF INDUSTRIAL SCIENCE AND TECHNOLOGY)  * column 13, line 8 - column 15, line 32; figure 5 *	1-2, 8-10, 13-15	
	---		
	---/---		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04 MARCH 1992	Examiner LE CORNEC N. D. R.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number

EP 91 11 6485

Page 2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	PATENT ABSTRACTS OF JAPAN vol. 12, no. 110 (C-486)(2957) 8 April 1988 & JP-A-62 236 485 ( AGENCY OF INDUSTRIAL SCIENCE AND TECHNOLOGY ) 16 October 1987 * abstract * & CHEMICAL ABSTRACTS, vol. 108, no. 25, 20 June 1988, Columbus, Ohio, US; abstract no. 217272, SAKAKI Y. ET AL: 'Chimeric rat liver cytochrome P-450 and its gene expression in yeast' page 163 ; column L ; * abstract *	1,3-5	
X	--- JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, no. 34, 1988, BALTIMORE US pages 18138 - 18142; KYUICHI OKUDA ET AL: 'Purification and characterization of 5beta-cholestane-3alpha,7alpha,12alpha-triol 27-hydroxylase from female rat liver mitochondria' * page 18139, left column, line 65 - right column, line 29; table III * * page 18142, left column, line 50 - line 59; figure 5 *	15	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
X	--- JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, no. 28, 1988, BALTIMORE US pages 14256 - 14260; OSAMU MASUMOTO ET AL: 'Purification and characterization of vitamin D 25-hydroxylase from rat liver mitochondria' * page 14257, left column, line 40 - line 74; tables II,III * --- -/--	15	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04 MARCH 1992	Examiner LE CORNEC N. D. R.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			

EPO FORM 1500 CL.52 (P0481)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number

EP 91 11 6485

Page 3

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,X	FEBS LETTERS. vol. 280, no. 2, March 1991, AMSTERDAM NL pages 367 - 370; MEGUMI AKIYOSHI-SHIBATA ET AL: 'Expression of rat liver vitamin D3 25-hydroxylase cDNA in Saccharomyces cerevisiae' * the whole document *	1-2, 9-10, 13-14,15	
P,X	FEBS LETTERS. vol. 274, no. 1,2, November 1990, AMSTERDAM NL pages 175 - 177; E. USUI ET AL: 'Unique property of liver mitochondrial P450 to catalyse the two physiologically important reactions involved in both cholesterol catabolism and vitamin D activation' * the whole document *	3,5,15	
A	CHEMICAL ABSTRACTS, vol. 108, no. 25, 20 June 1988, Columbus, Ohio, US; abstract no. 218092, DAHLBAECK H. ET AL: '25-hydroxylation of vitamin D3 by a cytochrome P450 from rabbit liver mitochondria.' page 240 ;column R ; * abstract * & BIOCHEM. J. vol. 252, no. 1, 1988, pages 207 - 213;	15	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Place of search THE HAGUE		Date of completion of the search 04 MARCH 1992	Examiner LE CORNEC N. D. R.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document			

EPO FORM 150 (03.92) (P0001)